



PPARgamma agonists can be expected to potentiate the efficacy of metronomic chemotherapy through CD36 up-regulation

Mark F. McCarty *, Jorge Barroso-Aranda, Francisco Contreras

Oasis of Hope Hospital, Tijuana, Mexico

Received 15 December 2006; accepted 18 December 2006

Summary The ability of metronomic chemotherapy to induce endothelial apoptosis has been traced to increased endothelial expression of thrombospondin-1, which activates endothelial CD36 receptors, triggering the extrinsic apoptotic pathway. Endothelial expression of CD36 is variable. Recent studies show that PPARgamma agonists – previously shown to have angiostatic activity – can markedly boost endothelial expression of CD36, thereby potentiating the apoptotic response of endothelial cells to thrombospondin-1-mimetic peptides. Thus, concurrent administration of PPARgamma agonists would be expected to enhance the efficacy of metronomic chemotherapy. These considerations may help to rationalize recent reports that a regimen consisting of low-dose trofosamide, pioglitazone, and a cox-2 inhibitor achieves tumor regression or prolonged tumor stasis in a meaningful proportion of cancer patients. The angiostatic efficacy of metronomic chemotherapy complemented by PPARgamma agonist administration would likely be potentiated by ancillary measures that block the survival signals evoked by endothelial growth factors such as VEGF or angiopoietin-1.

© 2007 Elsevier Ltd. All rights reserved.

Role of thrombospondin-1/CD36 interaction in metronomic chemotherapy

There is cogent evidence that increased production of thrombospondin 1 (TSP-1) is an obligate mediator of the endothelial apoptosis induced by metronomic therapy [1–3]. Thus, continual exposure of cultured endothelial cells to modest concentrations of genotoxic agents (too low to kill cancer

cells directly) increases their production and secretion of TSP-1, which can interact with the CD36 receptors typically expressed by endothelium to induce apoptosis [4–7]. The pro-apoptotic pathway triggered by CD36 activation appears to involve p59^{fyn}-mediated stimulation of c-Jun NH2-terminal kinase and p38 MAP kinase, resulting in increased Fas ligand expression. This ligand then interacts with Fas – which metronomic chemotherapy also induces [8,9] – to trigger extrinsic apoptosis [5,8–10]. A monoclonal antibody that inhibits TSP-1 binding to CD36 largely abrogates the adverse impact of continuous genotoxin exposure on the proliferation and survival of endothelial cells,

* Corresponding author. Address: Natural Alternatives International, 1185 Linda Vista Drive, San Marcos, CA 92078, United States.

E-mail address: mccarty@pantox.com (M.F. McCarty).

pointing to the TSP-1/CD36 interaction as key to the evoked apoptosis [1]. Moreover, metronomic chemotherapy with cyclophosphamide fails to impede tumor growth in mice that are TSP-1-null, whereas intermittent high-dose cyclophosphamide retains antitumor activity in these mice [1]. Why genotoxins provoke TSP-1 expression in endothelial cells remains unclear, though a likely possibility is induction via p53 activation [11]; p. 53 can likewise induce Fas expression [12].

PPARgamma agonists Up-regulate CD36

The extent of expression of CD36 by endothelial cells varies as a function of their location, and this expression is substantially diminished in cultured human microvascular endothelial cells after multiple passages. Recently, Huang and colleagues have established that exposure to various PPARgamma agonists, including the drugs troglitazone and rosiglitazone, markedly boosts CD36 expression in late passage endothelial cells, restoring their sensitivity to TSP-1 [13]. Analogous induction of CD36 by PPARgamma had previously been observed in various hepatic cells [14]. This finding is of particular interest in light of multiple reports that PPARgamma agonists exert marked anti-angiogenic tumor-retardant activity in vivo in rodent xenograft models [15,16]; Whether CD36 induction contributes importantly to this effect is not clear. In vitro, PPARgamma agonists decrease VEGF receptor expression by cultured endothelial cells, and reduce their responsiveness to exogenous growth factors [17]. This latter effect may reflect, in part, an up-regulation of PTEN that down-regulates Akt activation [18]. The anti-angiogenic impact of PPARgamma is not contingent on PPARgamma expression by the tumor, though in some tumors which do express this receptor PPARgamma agonists can decrease production of angiogenic factors [19]. In any case, treatment with PPARgamma agonists has been shown to potentiate the anti-angiogenic impact of concurrent administration of ABT510 [13], a modified peptide derived from TSP-1 that has high affinity for CD36 and mimics the pro-apoptotic effects of TSP-1 on endothelial cells [20,21]. In particular, the extent of apoptosis in tumor endothelial cells was markedly higher in animals treated jointly with rosiglitazone and ABT510 than in animals receiving either agent alone; the effect appeared to be synergistic rather than merely additive [13].

Whether ABT510 – currently being assessed in phase I studies [22,23] – or other TSP-1 mimetics, ever become available for clinical use remains to

be seen; they require daily parenteral administration. However, as noted above, it is feasible to evoke autocrine TSP-1 expression in endothelial cells through use of metronomic chemotherapy, which is an established clinical technique – a regimen involving low-dose cyclophosphamide and methotrexate having shown very worthwhile efficacy in late-stage breast cancer [24–26]. Thus, it is proposed that treatment with PPARgamma agonist drugs (such as glitazones) will potentiate the tumor-retardant activity of metronomic chemotherapy by boosting endothelial expression of CD36. In addition, PPARgamma-mediated down-regulation of Akt activity might be expected to potentiate the apoptotic response to metronomic chemotherapy, inasmuch as Akt works in a number of complementary ways – most notably, BAD phosphorylation – to oppose apoptosis in endothelial cells and other tissues [27–31]. Testing the interaction of metronomic chemotherapy and PPARgamma agonists would make good sense in any case, as both of these modalities have demonstrated good angiostatic tumor-retardant activity in rodent xenograft models.

Clinical experience

Indeed, Vogt and colleagues at the University of Regensburg have evaluated a combination angiostatic regimen – entailing daily administration of pioglitazone (45 mg), rofecoxib (25 mg) and trofosfamide (50 mg t.i.d.) – in patients with late-stage melanoma, soft-tissue sarcomas, and various vascular tumors (including kaposi sarcoma). A significant proportion of these patients achieved either objective responses or prolonged disease stabilization [32–35]. These findings evidently do not allow any conclusions regarding the contributions of the individual components of this regimen to the observed responses, but they are certainly consistent with the possibility that combining metronomic chemotherapy with PPARgamma agonists can achieve worthwhile angiostatic activity. (The inclusion of rofecoxib in the Regensburg regimen reflects the angiostatic utility of cox-2 inhibition; induction of cox-2 in plays a role in endothelial tube formation, pericyte recruitment, and endothelial cell survival during early angiogenesis [36,37]).

Adjunctive potentiators of metronomic chemotherapy

Of related interest is a study by Pietras and Hanahan demonstrating that concurrent inhibition

of VEGFR-2 and PDGF receptors greatly boosts the angiostatic efficacy of metronomic chemotherapy in transgenic mice bearing autogenous pancreatic islet tumors [38]. Presumably, the metronomic chemotherapy provides a TSP-1-mediated apoptotic stimulus to tumor endothelial cells, while inhibition of VEGF and PDGF activity eliminates crucial survival signals that protect endothelial cells from apoptosis. (The effect of PDGFR inhibition appears to be indirect – it prevents the interaction of free endothelial cells with pericytes that can provide survival signals such as angiopoietin-1 [39,40]). Other researchers have demonstrated synergism between metronomic vinblastine and an antibody targeting VEGFR-2 in a neuroblastoma xenograft model [41]. These findings are particularly exciting in light of the fact that the drug sunitinib, which can inhibit both VEGFR-2 and PDGFR in low nanomolar concentrations achievable with acceptable toxicity [42], has recently become available for clinical use in the US [43,44]. Thus, one can envision an angiostatic regimen combining metronomic chemotherapy, sunitinib, and a PPARgamma agonist – possibly in conjunction with other well-tolerated oral agents such as fish oil, cox-2 inhibitors, silibinin, and salicylate that also have angiostatic potential [45]. Salicylate and silibinin may be of particular interest in this regard, as the endothelial survival signals provided by VEGFR-2 and various integrins are contingent on activation of NF- κ B and its upstream activator IKK-2 [46–49] – which these agents target [50,51]; NF- κ B also induces the Tie2 receptor and plays a key role in capillary tube formation [52,53]. There seems to be a broad consensus that an angiostatic therapy that is both effective and clinically tolerable will require use of multiple agents – preferably orally administrable ones – that address multiple targets in a complementary fashion.

In passing, it should be noted that, while PPARgamma agonists appear likely to potentiate metronomic chemotherapy, they also have the potential to increase the responsiveness of certain tumors to standard chemotherapy. In tumors which express PPARgamma and still carry at least one functional PTEN gene, PPARgamma agonists could be expected to down-regulate Akt activity by boosting PTEN expression, just as they do in endothelial cells [54–60]. Increased PTEN expression in cancer cells has been shown to exert a chemo- and radiosensitizing effect [61–64]. Furthermore, long-term down-regulation of tumor Akt activity would likely have a favorable impact on tumor aggressiveness [65].

References

- [1] Bocci G, Francia G, Man S, Lawler J, Kerbel RS. Thrombospondin 1, a mediator of the antiangiogenic effects of low-dose metronomic chemotherapy. *Proc Natl Acad Sci USA* 2003;100:12917–22.
- [2] Ng SS, Figg WD. Upregulation of endogenous angiogenesis inhibitors: a mechanism of action of metronomic chemotherapy. *Cancer Biol Ther* 2004;3:1212–3.
- [3] Damber JE, Vallbo C, Albertsson P, Lennernas B, Norrby K. The anti-tumour effect of low-dose continuous chemotherapy may partly be mediated by thrombospondin. *Cancer Chemother Pharmacol* 2006;58:354–60.
- [4] Jimenez B, Volpert OV, Crawford SE, Febbraio M, Silverstein RL, Bouck N. Signals leading to apoptosis-dependent inhibition of neovascularization by thrombospondin-1. *Nat Med* 2000;6:41–8.
- [5] Jimenez B, Volpert OV, Reiher F, Chang L, Munoz A, Karin M, et al. c-Jun N-terminal kinase activation is required for the inhibition of neovascularization by thrombospondin-1. *Oncogene* 2001;20:3443–8.
- [6] Reiher FK, Volpert OV, Jimenez B, Crawford SE, Dinney CP, Henkin J, et al. Inhibition of tumor growth by systemic treatment with thrombospondin-1 peptide mimetics. *Int J Cancer* 2002;98:682–9.
- [7] Armstrong LC, Bornstein P. Thrombospondins 1 and 2 function as inhibitors of angiogenesis. *Matrix Biol* 2003;22:63–71.
- [8] Quesada AJ, Nelius T, Yap R, Zaichuk TA, Alfranca A, Filleur S, et al. In vivo upregulation of CD95 and CD95L causes synergistic inhibition of angiogenesis by TSP1 peptide and metronomic doxorubicin treatment. *Cell Death Differ* 2005;12:649–58.
- [9] Yap R, Veliceasa D, Emmenegger U, Kerbel RS, McKay LM, Henkin J, et al. Metronomic low-dose chemotherapy boosts CD95-dependent antiangiogenic effect of the thrombospondin peptide ABT-510: a complementation antiangiogenic strategy. *Clin Cancer Res* 2005;11:6678–85.
- [10] Jimenez B, Volpert OV, Crawford SE, Febbraio M, Silverstein RL, Bouck N. Signals leading to apoptosis-dependent inhibition of neovascularization by thrombospondin-1. *Nat Med* 2000;6:41–8.
- [11] Harada H, Nakagawa K, Saito M, Kohno S, Nagato S, Furukawa K, et al. Introduction of wild-type p53 enhances thrombospondin-1 expression in human glioma cells. *Cancer Lett* 2003;191:109–19.
- [12] Owen-Schaub LB, Zhang W, Cusack JC, Angelo LS, Santee SM, Fujiwara T, et al. Wild-type human p53 and a temperature-sensitive mutant induce Fas/APO-1 expression. *Mol Cell Biol* 1995;15:3032–40.
- [13] Huang H, Campbell SC, Bedford DF, Nelius T, Veliceasa D, Shroff EH, et al. Peroxisome proliferator-activated receptor gamma ligands improve the antitumor efficacy of thrombospondin peptide ABT510. *Mol Cancer Res* 2004;2:541–50.
- [14] Malerod L, Sporstol M, Juvet LK, Mousavi A, Gjoen T, Berg T. Hepatic scavenger receptor class B, type I is stimulated by peroxisome proliferator-activated receptor gamma and hepatocyte nuclear factor 4alpha. *Biochem Biophys Res Commun* 2003;305:557–65.
- [15] Panigrahy D, Singer S, Shen LQ, Butterfield CE, Freedman DA, Chen EJ, et al. PPARgamma ligands inhibit primary tumor growth and metastasis by inhibiting angiogenesis. *J Clin Invest* 2002;110:923–32.

- [16] Panigrahy D, Huang S, Kieran MW, Kaipainen A. PPARgamma as a Therapeutic Target for Tumor Angiogenesis and Metastasis. *Cancer Biol Ther* 2005;4:687–93.
- [17] Xin X, Yang S, Kowalski J, Gerritsen ME. Peroxisome proliferator-activated receptor gamma ligands are potent inhibitors of angiogenesis in vitro and in vivo. *J Biol Chem* 1999;274:9116–21.
- [18] Goetze S, Bungenstock A, Czupalla C, Eilers F, Stawowy P, Kintscher U, et al. Leptin induces endothelial cell migration through Akt, which is inhibited by PPARgamma-ligands. *Hypertension* 2002;40:748–54.
- [19] Keshamouni VG, Arenberg DA, Reddy RC, Newstead MJ, Anthwal S, Standiford TJ. PPAR-gamma activation inhibits angiogenesis by blocking ELR + CXC chemokine production in non-small cell lung cancer. *Neoplasia* 2005;7:294–301.
- [20] Dawson DW, Volpert OV, Pearce SF, Schneider AJ, Silverstein RL, Henkin J, et al. Three distinct D-amino acid substitutions confer potent antiangiogenic activity on an inactive peptide derived from a thrombospondin-1 type 1 repeat. *Mol Pharmacol* 1999;55:332–8.
- [21] Reiher FK, Volpert OV, Jimenez B, Crawford SE, Dinney CP, Henkin J, et al. Inhibition of tumor growth by systemic treatment with thrombospondin-1 peptide mimetics. *Int J Cancer* 2002;98:682–9.
- [22] Hoekstra R, de Vos FY, Eskens FA, Gietema JA, van der GA, Groen HJ, et al. Phase I safety, pharmacokinetic, and pharmacodynamic study of the thrombospondin-1-mimetic angiogenesis inhibitor ABT-510 in patients with advanced cancer. *J Clin Oncol* 2005;23:5188–97.
- [23] Hoekstra R, de Vos FY, Eskens FA, de Vries EG, Uges DR, Knight R, et al. Phase I study of the thrombospondin-1-mimetic angiogenesis inhibitor ABT-510 with 5-fluorouracil and leucovorin: a safe combination. *Eur J Cancer* 2006;42:467–72.
- [24] Bocci G, Tuccori M, Emmenegger U, Liguori V, Falcone A, Kerbel RS, et al. Cyclophosphamide–methotrexate ‘metronomic’ chemotherapy for the palliative treatment of metastatic breast cancer. A comparative pharmacoeconomic evaluation. *Ann Oncol* 2005;16:1243–52.
- [25] Colleoni M, Orlando L, Sanna G, Rocca A, Maisonneuve P, Peruzzotti G, et al. Metronomic low-dose oral cyclophosphamide and methotrexate plus or minus thalidomide in metastatic breast cancer: antitumor activity and biological effects. *Ann Oncol* 2006;17:232–8.
- [26] Orlando L, Cardillo A, Rocca A, Balduzzi A, Ghisini R, Peruzzotti G, et al. Prolonged clinical benefit with metronomic chemotherapy in patients with metastatic breast cancer. *Anticancer Drugs* 2006;17:961–7.
- [27] Roos WP, Kaina B. DNA damage-induced cell death by apoptosis. *Trends Mol Med* 2006;12:440–50.
- [28] Liu XM, Ensenat D, Wang H, Schafer AI, Durante W. Physiologic cyclic stretch inhibits apoptosis in vascular endothelium. *FEBS Lett* 2003;541:52–6.
- [29] Nishimura K, Li W, Hoshino Y, Kadohama T, Asada H, Ohgi S, et al. Role of AKT in cyclic strain-induced endothelial cell proliferation and survival. *Am J Physiol Cell Physiol* 2006;290:C812–21.
- [30] Ohi N, Nishikawa Y, Tokairin T, Yamamoto Y, Doi Y, Omori Y, et al. Maintenance of Bad phosphorylation prevents apoptosis of rat hepatic sinusoidal endothelial cells in vitro and in vivo. *Am J Pathol* 2006;168:1097–106.
- [31] Huang J, Kontos CD. PTEN modulates vascular endothelial growth factor-mediated signaling and angiogenic effects. *J Biol Chem* 2002;277:10760–6.
- [32] Vogt T, Hafner C, Bross K, Bataille F, Jauch KW, Berand A, et al. Antiangiogenic therapy with pioglitazone, rofecoxib, and metronomic trofosamide in patients with advanced malignant vascular tumors. *Cancer* 2003;98:2251–6.
- [33] Reichle A, Bross K, Vogt T, Bataille F, Wild P, Berand A, et al. Pioglitazone and rofecoxib combined with angiostatically scheduled trofosamide in the treatment of far-advanced melanoma and soft tissue sarcoma. *Cancer* 2004;101:2247–56.
- [34] Coras B, Hafner C, Reichle A, Hohenleutner U, Szeimies RM, Landthaler M, et al. Antiangiogenic therapy with pioglitazone, rofecoxib, and trofosamide in a patient with endemic kaposi sarcoma. *Arch Dermatol* 2004;140:1504–7.
- [35] Reichle A, Vogt T, Kunz-Schughart L, Bretschneider T, Bachthaler M, Bross K, et al. Anti-inflammatory and angiostatic therapy in chemorefractory multisystem Langerhans’ cell histiocytosis of adults. *Br J Haematol* 2005;128:730–2.
- [36] Gately S, Li WW. Multiple roles of COX-2 in tumor angiogenesis: a target for antiangiogenic therapy. *Semin Oncol* 2004;31:2–11.
- [37] Monnier Y, Zaric J, Ruegg C. Inhibition of angiogenesis by non-steroidal anti-inflammatory drugs: from the bench to the bedside and back. *Curr Drug Targets Inflamm Allergy* 2005;4:31–8.
- [38] Pietras K, Hanahan D. A multitargeted, metronomic, and maximum-tolerated dose ‘chemo-switch’ regimen is antiangiogenic, producing objective responses and survival benefit in a mouse model of cancer. *J Clin Oncol* 2005;23:939–52.
- [39] Bergers G, Song S, Meyer-Morse N, Bergsland E, Hanahan D. Benefits of targeting both pericytes and endothelial cells in the tumor vasculature with kinase inhibitors. *J Clin Invest* 2003;111:1287–95.
- [40] Erber R, Thurnher A, Katsen AD, Groth G, Kerger H, Hammes HP, et al. Combined inhibition of VEGF and PDGF signaling enforces tumor vessel regression by interfering with pericyte-mediated endothelial cell survival mechanisms. *FASEB J* 2004;18:338–40.
- [41] Klement G, Baruchel S, Rak J, Man S, Clark K, Hicklin DJ, et al. Continuous low-dose therapy with vinblastine and VEGF receptor-2 antibody induces sustained tumor regression without overt toxicity. *J Clin Invest* 2000;105:R15–24.
- [42] Faivre S, Delbaldo C, Vera K, Robert C, Lozahic S, Lassau N, et al. Safety, pharmacokinetic, and antitumor activity of SU11248, a novel oral multitarget tyrosine kinase inhibitor, in patients with cancer. *J Clin Oncol* 2006;24:25–35.
- [43] Motzer RJ, Hoosen S, Bello CL, Christensen JG. Sunitinib malate for the treatment of solid tumours: a review of current clinical data. *Expert Opin Invest Drugs* 2006;15:553–61.
- [44] Cabebe E, Wakelee H. Sunitinib: a newly approved small-molecule inhibitor of angiogenesis. *Drugs Today (Barc)* 2006;42:387–98.
- [45] McCarty MF, Block KI. Multifocal angiostatic therapy: an update. *Integr Cancer Ther* 2005;4:301–14.
- [46] Grosjean J, Kiriakidis S, Reilly K, Feldmann M, Paleolog E. Vascular endothelial growth factor signalling in endothelial cell survival: a role for NFkappaB. *Biochem Biophys Res Commun* 2006;340:984–94.
- [47] Scatena M, Almeida M, Chaisson ML, Fausto N, Nicosia RF, Giachelli CM. NF-kappaB mediates alphavbeta3 integrin-induced endothelial cell survival. *J Cell Biol* 1998;141:1083–93.
- [48] Rice J, Courter DL, Giachelli CM, Scatena M. Molecular mediators of alpha(v)beta(3)-induced endothelial cell survival. *J Vasc Res* 2006;43:422–36.
- [49] Klein S, de Fougerolles AR, Blaikie P, Khan L, Pepe A, Green CD, et al. Alpha 5 beta 1 integrin activates an NF-kappa B-dependent program of gene expression important for

- angiogenesis and inflammation. *Mol Cell Biol* 2002;22:5912–22.
- [50] Yin MJ, Yamamoto Y, Gaynor RB. The anti-inflammatory agents aspirin and salicylate inhibit the activity of I(kappa)B kinase-beta. *Nature* 1998;396:77–80.
- [51] Yoo HG, Jung SN, Hwang YS, Park JS, Kim MH, Jeong M, et al. Involvement of NF-kappaB and caspases in silibinin-induced apoptosis of endothelial cells. *Int J Mol Med* 2004;13:81–6.
- [52] DeBusk LM, Chen Y, Nishishita T, Chen J, Thomas JW, Lin PC. Tie2 receptor tyrosine kinase, a major mediator of tumor necrosis factor alpha-induced angiogenesis in rheumatoid arthritis. *Arthritis Rheum* 2003;48:2461–71.
- [53] Oitzinger W, Hofer-Warbinek R, Schmid JA, Koshelnick Y, Binder BR, De Martin R. Adenovirus-mediated expression of a mutant I(kappa)B kinase 2 inhibits the response of endothelial cells to inflammatory stimuli. *Blood* 2001;97:1611–7.
- [54] Patel L, Pass I, Coxon P, Downes CP, Smith SA, Macphee CH. Tumor suppressor and anti-inflammatory actions of PPAR-gamma agonists are mediated via upregulation of PTEN. *Curr Biol* 2001;11:764–8.
- [55] Farrow B, Evers BM. Activation of PPARgamma increases PTEN expression in pancreatic cancer cells. *Biochem Biophys Res Commun* 2003;301:50–3.
- [56] Chen WC, Lin MS, Bai X. Induction of apoptosis in colorectal cancer cells by peroxisome proliferators-activated receptor gamma activation up-regulating PTEN and inhibiting PI3K activity. *Chin Med J (Engl)* 2005;118:1477–81.
- [57] Lee SY, Hur GY, Jung KH, Jung HC, Lee SY, Kim JH, et al. PPAR-gamma agonist increase gefitinib's antitumor activity through PTEN expression. *Lung Cancer* 2006;51:297–301.
- [58] Teresi RE, Shaiu CW, Chen CS, Chatterjee VK, Waite KA, Eng C. Increased PTEN expression due to transcriptional activation of PPARgamma by Lovastatin and Rosiglitazone. *Int J Cancer* 2006;118:2390–8.
- [59] Han S, Roman J. Rosiglitazone suppresses human lung carcinoma cell growth through PPARgamma-dependent and PPARgamma-independent signal pathways. *Mol Cancer Ther* 2006;5:430–7.
- [60] Kim KY, Kim SS, Cheon HG. Differential anti-proliferative actions of peroxisome proliferator-activated receptor-gamma agonists in MCF-7 breast cancer cells. *Biochem Pharmacol* 2006;72:530–40.
- [61] Wick W, Furnari FB, Naumann U, Cavenee WK, Weller M. PTEN gene transfer in human malignant glioma: sensitization to irradiation and CD95L-induced apoptosis. *Oncogene* 1999;18:3936–43.
- [62] Tanaka M, Koul D, Davies MA, Liebert M, Steck PA, Grossman HB. MMAC1/PTEN inhibits cell growth and induces chemosensitivity to doxorubicin in human bladder cancer cells. *Oncogene* 2000;19:5406–12.
- [63] Huang H, Cheville JC, Pan Y, Roche PC, Schmidt LJ, Tindall DJ. PTEN induces chemosensitivity in PTEN-mutated prostate cancer cells by suppression of Bcl-2 expression. *J Biol Chem* 2001;276:38830–6.
- [64] Oki E, Baba H, Tokunaga E, Nakamura T, Ueda N, Futatsugi M, et al. Akt phosphorylation associates with LOH of PTEN and leads to chemoresistance for gastric cancer. *Int J Cancer* 2005;117:376–80.
- [65] Mitsiades CS, Mitsiades N, Koutsilieris M. The Akt pathway: molecular targets for anti-cancer drug development. *Curr Cancer Drug Targets* 2004;4:235–56.

Available online at www.sciencedirect.com

